

THE REGULARIZED TRACE FORMULA FOR THE STURM-LIOUVILLE EQUATION WITH SPECTRAL PARAMETER IN THE BOUNDARY CONDITIONS

NAMIG J. GULIYEV

ABSTRACT. The regularized trace formula of first order for the Sturm–Liouville equation with spectral parameter in the boundary conditions is obtained.

Consider the boundary value problem

$$-y''(x) + q(x)y(x) = \lambda y(x), \quad x \in [0, \pi], \quad (1)$$

$$\lambda(y'(0) - hy(0)) = h_1y'(0) - h_2y(0), \quad (2)$$

$$\lambda(y'(\pi) + Hy(\pi)) = H_1y'(\pi) + H_2y(\pi), \quad (3)$$

where $q(x) \in \mathcal{W}_2^1[0, \pi]$ is a real-valued function, $h, h_1, h_2, H, H_1, H_2 \in \mathbb{R}$ and

$$\delta := hh_1 - h_2 > 0, \quad \rho := HH_1 - H_2 > 0.$$

Let $\varphi(x, \lambda)$ be the solution of (1) satisfying the initial conditions

$$\varphi(0, \lambda) = h_1 - \lambda, \quad \varphi'(0, \lambda) = h_2 - \lambda h.$$

We denote $s := \sqrt{\lambda}$. Since $\varphi(x, \lambda)$ satisfies the integral equation

$$\varphi(x, \lambda) = (h_1 - s^2) \cos sx + (h_2 - hs^2) \frac{\sin sx}{s} + \int_0^x \frac{\sin[s(x-t)]}{s} q(t) \varphi(t, \lambda) dt$$

and the estimate

$$\varphi(x, \lambda) = O\left(|s|^2 e^{|\operatorname{Im} s|x}\right),$$

one can obtain the following representations:

$$\varphi(x, \lambda) = -s^2 \cos sx - \left(h + \frac{1}{2} \int_0^x q(t) dt\right) s \sin sx + O\left(e^{|\operatorname{Im} s|x}\right), \quad (4)$$

$$\begin{aligned} \varphi'(x, \lambda) = & s^3 \sin sx - \left(h + \frac{1}{2} \int_0^x q(t) dt\right) s^2 \cos sx - \\ & - \left(h_1 + \frac{q(x) + q(0)}{4} + \frac{h}{2} \int_0^x q(t) dt + \frac{1}{8} \left(\int_0^x q(t) dt\right)^2\right) s \sin sx - \\ & - \frac{s}{4} \int_0^x q'(t) \sin[s(x-2t)] dt + O\left(e^{|\operatorname{Im} s|x}\right). \quad (5) \end{aligned}$$

Denote by $\chi(\lambda)$ the characteristic function of (1)–(3):

$$\chi(\lambda) := (\lambda - H_1)\varphi'(\pi, \lambda) + (\lambda H - H_2)\varphi(\pi, \lambda).$$

From (4) and (5) we have

$$\begin{aligned} \chi(\lambda) := s^5 \sin s\pi - \omega s^4 \cos s\pi - \omega_1 s^3 \sin s\pi - \frac{s^3}{4} \int_0^\pi q'(t) \sin[s(\pi - 2t)] dt + \\ + O(|s|^2 e^{|\operatorname{Im} s|\pi}), \end{aligned} \quad (6)$$

where

$$\begin{aligned} \omega &:= h + H + \frac{1}{2} \int_0^\pi q(t) dt, \\ \omega_1 &:= h_1 + H_1 + hH + \frac{q(\pi) + q(0)}{4} + \frac{h + H}{2} \int_0^\pi q(t) dt + \frac{1}{8} \left(\int_0^\pi q(t) dt \right)^2. \end{aligned}$$

The eigenvalues of the problem (1)–(3) are real and coincide with the zeros of the function $\chi(\lambda)$. From (6) we obtain

$$s_n = n - 2 + \frac{\omega}{\pi(n-2)} + \frac{\zeta_n}{n^2}, \quad \{\zeta_n\} \in l_2.$$

Then

$$\lambda_n = s_n^2 = (n-2)^2 + \frac{2\omega}{\pi} + \frac{\xi_n}{n}, \quad \{\xi_n\} \in l_2. \quad (7)$$

Therefore

$$s_\lambda := \lambda_0 + \lambda_1 + \sum_{n=2}^\infty \left(\lambda_n - (n-2)^2 - \frac{2\omega}{\pi} \right) < \infty. \quad (8)$$

The series (8) is called *the regularized trace of first order* (or *the first regularized trace*) for the problem (1)–(3). Analogously, one can define *the second, the third* etc. regularized traces. Note that the second regularized trace formula for the Sturm–Liouville problem with spectral parameter in a boundary condition has been found in [1].

Since $\chi(\lambda)$ is an entire function of order $1/2$, from Hadamard's theorem (see [2, Section 4.2]), using (6) we obtain

$$\chi(\lambda) = \pi(\lambda - \lambda_0)(\lambda - \lambda_1)(\lambda - \lambda_2) \prod_{n=3}^\infty \frac{\lambda_n - \lambda}{(n-2)^2} \quad (9)$$

Let $\lambda = -\mu^2$. We calculate the sum s_λ of the series (8) by comparing the asymptotic expressions obtained from formulas (6) and (9) as $\mu \rightarrow \infty$. From the formula (9) we have

$$\begin{aligned} \chi(-\mu^2) &= -\pi(\mu^2 + \lambda_0)(\mu^2 + \lambda_1)(\mu^2 + \lambda_2) \prod_{n=3}^\infty \frac{\lambda_n + \mu^2}{(n-2)^2} = \\ &= -\frac{(\mu^2 + \lambda_0)(\mu^2 + \lambda_1)(\mu^2 + \lambda_2) \operatorname{sh} \pi \mu}{\mu} \Phi(\mu), \end{aligned} \quad (10)$$

where

$$\Phi(\mu) = \prod_{n=1}^{\infty} \left(1 - \frac{n^2 - \lambda_{n+2}}{\mu^2 + n^2} \right).$$

We need the formulas (see [3])

$$\sum_{k=2}^{\infty} \frac{1}{k} \sum_{n=1}^{\infty} \frac{|n^2 - \lambda_{n+2}|^k}{(\mu^2 + n^2)^k} = o\left(\frac{1}{\mu^2}\right) \quad (11)$$

and

$$\sum_{n=1}^{\infty} \frac{1}{\mu^2 + n^2} = \frac{\pi}{2\mu} - \frac{1}{2\mu^2} + o\left(\frac{1}{\mu^2}\right). \quad (12)$$

Using (7) and (12) we have

$$\begin{aligned} \frac{1}{\mu^2} \sum_{n=1}^{\infty} \left(\lambda_{n+2} - n^2 - \frac{2\omega}{\pi} \right) \frac{n^2}{\mu^2 + n^2} &= \frac{1}{\mu^2} \sum_{n=1}^{\infty} \frac{n \xi_{n+2}}{\mu^2 + n^2} \leq \\ &\leq \frac{1}{\mu^2} \sqrt{\sum_{n=3}^{\infty} \xi_n^2} \sqrt{\sum_{n=1}^{\infty} \frac{n^2}{(\mu^2 + n^2)^2}} \leq \frac{\Omega}{\mu^2} \sqrt{\sum_{n=1}^{\infty} \frac{\mu^2 + n^2}{(\mu^2 + n^2)^2}} = \\ &= \frac{\Omega}{\mu^2} \sqrt{\sum_{n=1}^{\infty} \frac{1}{\mu^2 + n^2}} = o\left(\frac{1}{\mu^2}\right), \quad (13) \end{aligned}$$

where

$$\Omega := \sqrt{\sum_{n=3}^{\infty} \xi_n^2} < \infty.$$

From (11), (12) and (13) we calculate:

$$\begin{aligned} \ln \Phi(\mu) &= \sum_{n=1}^{\infty} \left(1 - \frac{n^2 - \lambda_{n+2}}{\mu^2 + n^2} \right) = - \sum_{n=1}^{\infty} \sum_{k=1}^{\infty} \frac{1}{k} \left(\frac{n^2 - \lambda_{n+2}}{\mu^2 + n^2} \right)^k = \\ &= - \sum_{k=1}^{\infty} \frac{1}{k} \sum_{n=1}^{\infty} \left(\frac{n^2 - \lambda_{n+2}}{\mu^2 + n^2} \right)^k = - \sum_{n=1}^{\infty} \frac{n^2 - \lambda_{n+2}}{\mu^2 + n^2} + o\left(\frac{1}{\mu^2}\right) = \\ &= \frac{2\omega}{\pi} \sum_{n=1}^{\infty} \frac{1}{\mu^2 + n^2} + \sum_{n=1}^{\infty} \left(\lambda_{n+2} - n^2 - \frac{2\omega}{\pi} \right) \frac{1}{\mu^2 + n^2} + o\left(\frac{1}{\mu^2}\right) = \frac{\omega}{\mu} - \frac{\omega}{\pi\mu^2} + \\ &+ \frac{1}{\mu^2} \sum_{n=1}^{\infty} \left(\lambda_{n+2} - n^2 - \frac{2\omega}{\pi} \right) - \frac{1}{\mu^2} \sum_{n=1}^{\infty} \left(\lambda_{n+2} - n^2 - \frac{2\omega}{\pi} \right) \frac{n^2}{\mu^2 + n^2} + \\ &+ o\left(\frac{1}{\mu^2}\right) = \frac{\omega}{\mu} - \frac{\omega}{\pi\mu^2} + \frac{1}{\mu^2} \left(s_{\lambda} - \lambda_0 - \lambda_1 - \lambda_2 + \frac{2\omega}{\pi} \right) + o\left(\frac{1}{\mu^2}\right) = \\ &= \frac{\omega}{\mu} + \frac{1}{\mu^2} \left(s_{\lambda} - \lambda_0 - \lambda_1 - \lambda_2 + \frac{\omega}{\pi} \right) + o\left(\frac{1}{\mu^2}\right). \end{aligned}$$

Therefore

$$\Phi(\mu) = 1 + \frac{\omega}{\mu} + \frac{1}{\mu^2} \left(s_{\lambda} - \lambda_0 - \lambda_1 - \lambda_2 + \frac{\omega}{\pi} + \frac{\omega^2}{2} \right) + o\left(\frac{1}{\mu^2}\right)$$

and from (10)

$$\chi(-\mu^2) = \frac{e^{\pi\mu}}{2} \left(-\mu^5 - \omega\mu^4 + \mu^3 \left(s_\lambda + \frac{\omega}{\pi} + \frac{\omega^2}{2} \right) + o(\mu^3) \right). \quad (14)$$

The estimate (6) implies

$$\chi(-\mu^2) = \frac{e^{\pi\mu}}{2} (-\mu^5 - \omega\mu^4 + \omega_1\mu^3 + o(\mu^3)). \quad (15)$$

Comparing (14) and (15) we obtain the following trace formula:

$$\begin{aligned} \lambda_0 + \lambda_1 + \sum_{n=2}^{\infty} \left(\lambda_n - (n-2)^2 - \frac{2\omega}{\pi} \right) = \\ = h_1 + H_1 + \frac{q(\pi) + q(0)}{4} - \frac{h + H}{\pi} - \frac{h^2 + H^2}{2} - \frac{1}{2\pi} \int_0^{\pi} q(t) dt. \end{aligned}$$

REFERENCES

- [1] Albayrak I., Bayramoğlu M. and Adıgüzelov E. *The second regularized trace formula for the Sturm–Liouville problem with spectral parameter in a boundary condition*. Methods Funct. Anal. Topology, 2000, v.6, No.3, pp.1–8.
- [2] Levin B.Ya. *Lectures on entire functions*. Providence, RI: Amer. Math. Soc., 1996.
- [3] Levitan B.M. *Calculation of the regularized trace for the Sturm–Liouville operator*. Uspekhi Mat. Nauk, 1964, v.19, No.1(115), pp.161–165. (Russian)

Namig J. Guliyev

Institute of Mathematics and Mechanics of NAS of Azerbaijan

9 F.Agayev str., AZ1141, Baku, Azerbaijan.

E-mail: njguliyev@yahoo.com